Structure and vibrational spectrum of isopropenyl radical, $H_2C–C^{\bullet}=CH_2$: ab initio MO study and normal coordinate analysis
V.G. Avakyan (Moscow, U.S.S.R.) 99
Temperature- and solvent-dependent Raman bands of n-propylamine-*N*-d_2: a conformational study
L.A.E. Batista de Carvalho, A.M. Amorim da Costa and J.J.C. Teixeira-Dias (Coimbra, Portugal) 105
The infrared spectra of methyl(*O*-methyl)dithiophosphonic acid. Multiplicity of some bands due to different conformers
R. Grecu, R. Constantinescu, I. Silaghi-Dumitrescu and I. Haiduc (Cluj-Napoca, Roumania) . 111
High overtones investigation of non-equivalent C–H bonds in pyridine and 2,6-lutidine by thermal lensing spectroscopy
R. Bini, P. Foggi (Firenze, Italy) and R.G. Della Valle (Venice, Italy) 117
Infrared and Raman study of some heteropolyacid hydrates
U. Mioč (Belgrade, Yugoslavia), Ph. Colomban (Palaiseau, France) and A. Novak (Thiais, France) 123
Influence of the conjugation effect on the UV-spectra of 4-substituted pyridine *N*-oxides and their conjugated acids
L. Chmurzyński and A. Liwo (Gdańsk, Poland) 129
Investigation of bis-crown ethers for ion-selective electrodes by FT-IR spectroscopy
O. Egyed, V.P. Izvekov, K. Toth, S. Holly and E. Pungor (Budapest, Hungary) 135
IR study of the cholesteryl n-alkanoates conformation mobility
G.A. Puchkovskaya (Kiev, U.S.S.R.) and A.A. Yakubov (Samarkand, U.S.S.R.) 141
Raman spectroscopy and conformations of 1,3-dithia-5,6-benzocycloheptenes
P.N. Sobolev, I.Kh. Shakirov and R.R. Shgidullin (Kazan, U.S.S.R.) 147
IR- and NMR-spectra and structure of acyl salts of azomethins
C.Ju. Chotiy, M.S. Platitsa, V.I. Rybachenko, V.F. Galat and E.V. Titov (Donetsk, U.S.S.R.) 153
FTIR-CD of induced cholesteric solutions
D. Tsankov, B. Jordanov, D. Kolev (Sofia, Bulgaria) and E.H. Korte (Dortmund, F.R.G.) 159
Hindered rotation in amidines
I. Wawer (Warsaw, Poland) 165
On the "selenation" method of assignment of organic sulphur compounds vibrational spectra
O.M. Fugaeva, A.V. Zibarev, I.K. Korobeinicheva and G.G. Furin (Novosibirsk, U.S.S.R.) ... 169
STO-4G–SCF–MO calculations and NMR measurements for alkenyl phosphonic acids
Y.G. Smeyers, A. Hernandez-Laguna, C.I. Sainz-Diaz, M. Fernandez-Ibañez, E. Galvez-Ruano and M.S. Arias-Perez (Madrid, Spain) 175
Vibrational spectra and the structure of I and II group metals caprolactamates
I.A. Garbuzova, L.A. Checulaeva, L.B. Danilevskaya, V.A. Kotelnikov, V.V. Kurashev, V.V. Gavrilenko and B.V. Lokshin (Moscow, U.S.S.R.) 183
IR and Raman investigations on 5-element-1-stanna-bicyclo[3.3.$0^{1,5}$]octanes and 2,8-dithia-5-element-1-stanna-bicyclo[3.3.0^{15}]-octanes
K. Schenzel, A. Kolbe (Halle, G.D.R.), A. Sokołovska (Warsaw, Poland) and P. Reich (Berlin, G.D.R.) 189
Photoelectron spectra, orbital interaction, and conformations of alkylalkenylcyclopropanes and alkylalkenylcyclopropenes
V.V. Plemenkov, O.Ju. Butenko, V.V. Zverev, L.V. Ermolaeva, V.M. Vakar (Kazan, U.S.S.R.), A.V. Ignatchenko and I.G. Bolesov (Moscow, U.S.S.R.) 195
UPS investigation of 1-substituted pyrroles
L. Nyulászi, T. Tóth, Gy. Zsombok, G. Csonka, J. Réffy, J. Nagy and T. Veszprémi (Budapest, Hungary) 201
Molecular structure and conformation changes of mesogen crystals
L.M. Babkov (Saratov, U.S.S.R.), N.A. Golovina, A.V. Kosov, A.P. Polishchuk and G.A. Puchkovskaya (Kiev, U.S.S.R.) 207

IR spectra of ethyl acetate and perfluoro-tert-butanol–ethyl acetate complex at different temperatures
M.I. Redondo, M.V. Garcia, M.A. Raso (Madrid, Spain) and W.A.P. Luck (Marburg, F.R.G.) 213

MOLECULAR CONSTANTS
Analysis of the three lowest frequency vibration–rotation bands of diazomethane
L. Nemes (Budapest, Hungary), J. Vogt (Ulm, F.R.G.) and M. Winnewisser (Giessen, F.R.G.) 219
Laser-induced fluorescence spectroscopy of SiF produced by IR multiple-photon dissociation of SiF_4
V.K. Popov (Troitsk, U.S.S.R.), M. Rossberg, W. Strube, J. Wollbrandt and E. Linke (Berlin, G.D.R.) 225
On application of vibrational spectroscopy to the study of chemical bonds in complex molecules
E.N. Yurchenko (Leningrad, U.S.S.R.), N.N. Bulgakov and G.E. Osipova (Novosibirsk, U.S.S.R.) 231
Vibrational spectra of monothiocarbamates. I. Assignment and tentative force field of *S*-methyl-*N*,*N*-dimethylthiocarbamate
G. Keresztury, G. Besenyei and S. Holly (Budapest, Hungary) 237
Interpretation of vibrational spectra by intensity calculations with semi-empirical methods
H. Mikosch (Vienna, Austria) 243
Band intensities in vibrational spectra and ester structure
V.L. Furer (Kazan, U.S.S.R.) 249
Charge separation study in bisbenzene complexes of metals by theoretical analysis of infrared intensities
I.I. Greenwald and B.V. Lokshin (Moscow, U.S.S.R.) 255
Internal vibrational energy redistribution and vibrationally induced nonlinearity of HCN
W. Quapp (Leipzig, G.D.R.) 261
Electronic absorption spectra of chromyl compounds
M. Sowińska, J. Myrczek and A. Bartecki (Wrocław, Poland) 267
Fluorescence analysis of laser produced radicals and atoms
H. Hohmann and W. Radloff (Berlin, G.D.R.) 273
Modelling of temperature influence on the non-linear vibronic spectra of polyatomic molecules
P.P. Kircheva and S.D. Simeonov (Sofia, Bulgaria) 279
Interpretation of infrared intensities based on the atomic polar tensor partition. The ROH molecules (R=H, CH_3, NH_2, OH, F)
M.M.C. Ferreira (Campinas, Brazil) 285
Application of the Pade-form Hamiltonians for processing of vibration–rotation spectra of diatomic and triatomic molecules
V.F. Golovko, S.N. Mikhailenko and VI.G. Tyuterev (Tomsk, U.S.S.R.) 291
High pressure Raman study of Fermi resonance spectrum in gaseous carbon dioxide
A. Hacura, A. Bródka, L. Nikiel (Katowice, Poland) and F.G. Baglin (Reno, NV, U.S.A.) 297
The planar harmonic force field for 1,2-carbodicyanocyclobuten-3,4-dione dianion
B. Lunelli and M.G. Giorgini (Bologna, Italy) 303
FTIR linear dichroic spectra of substances dissolved and oriented in a nematic liquid crystal
M. Belhakem and B. Jordanov (Sofia, Bulgaria) 309
Utilization of overlay and ab initio force fields in the construction of empirical potential energy functions for conjugated molecules
L.-O. Pietilä, B. Mannfors (Helsinki, Finland) and K. Palmö (Ann Arbor, MI, U.S.A.) 315
MOLVIB — a flexible program for force field calculations
T. Sundius (Helsinki, Finland) 321
A new treatment of the ring puckering motions by periodic functions in the Hamiltonian
L. Sztraka (Budapest, Hungary) 327

The pressure dependence of the OH anharmonicity constant of perfluoro-tert-butanol in non-polar solvents
W.A.P. Luck and T.F. Mentel (Marburg, F.R.G.) 333
Effects of crossings and quasi-crossings of quasi-energy levels in spectra of molecules placed in a strong resonant field
A.A. Kiselev and A.V. Liapzev (Leningrad, U.S.S.R.) 339

SURFACE PHENOMENA
Surface plasmon resonance as a bioanalytical tool
R.P.H. Kooyman, H.E. de Bruijn, R.G. Eenink and J. Greve (Enschede, The Netherlands) 345
Selective spectroscopic characterization of Langmuir–Blodgett monolayers using SERRS
D. Battisti and R. Aroca (Windsor, Ont., Canada) 351
Application of SERS in photographic sciences
K. Kneipp (Berlin, G.D.R.) 357
IR spectra of hydrogen adsorbed on solids
G. Busca (Genoa, Italy) 363
UV–vis spectroscopic and theoretical studies about carbocation formation on solid surfaces
I. Kiricsi, Gy. Tasi (Szeged, Hungary), H. Förster (Hamburg, F.R.G.) and P. Fejes (Szeged, Hungary) 369
Electronic aspects of polyacene adsorptional complexes formation on silica surface and their fluorescence spectra
A.M. Yeremenko (Kiev, U.S.S.R.), L.A. Bykovskaya (Moscow, U.S.S.R.) and A.A. Chuiko (Kiev, U.S.S.R) 375
Influences of surface reflections on diffuse reflectance measurements
E.H. Korte and H. Staat (Dortmund, F.R.G.) 381
Optical high resolution study of quinizarin adsorbed on γ-alumine powder: indication of low frequency adsorbate motions
T. Basche' and C. Bräuchle (Munich, F.R.G.) 387
Sorption states of isoelectronic linear molecules in zeolites
H. Böse and H. Förster (Hamburg, F.R.G.) 393
Normal coordinate analysis of *trans*- and *cis*-butane adsorbed in zeolites A
H. Förster (Hamburg, F.R.G.) and O. Zakharieva-Pencheve (Sofia, Bulgaria) 399
Study of the geometry of cyclopropane adsorbed on a calcium ion in zeolites A by SCC-Xα calculations
O. Zakharieva-Pencheva (Sofia, Bulgaria), M. Grodzicki (New Orleans, LA, U.S.A.), H. Böse and H. Förster (Hamburg, F.R.G.) 405
IR and optical spectroscopy of structural changes of WO_3 electrochromic thin films
T.A. Gavrilko, V.I. Stepkin and I.V. Shiyanovskaya (Kiev, U.S.S.R.) 411
FI-IR study of the Brønsted acidity of phosphated and sulphated silica catalysts
M. Lion, M. Maache, J.C. Lavalley (Caen, France), G. Ramis, G. Busca, P.F. Rossi and V. Lorenzelli (Genoa, Italy) 417
Surface enhanced Raman scattering of benzo[*c*]cinnoline
N. Neto (Potenza, Italy), M. Nocentini, M. Muniz-Miranda and G. Sbrana (Firenze, Italy) 423
Salt-induced orientation of diazanaphthalenes adsorbed on silver sols, as detected by SERS
G. Sbrana, M. Muniz-Miranda (Firenze, Italy), N. Neto (Potenza, Italy) and M. Nocentini (Firenze, Italy) 429
Luminescence and transient absorption studies on bonding and mobility of nitrogen heterocycles on silica and alumina surfaces
D. Oelkrug, S. Uhl, M. Gregor, R. Lege (Tübingen, F.R.G.), G. Kelly and F. Wilkinson (Loughborough, Gt. Britain) 435
Vibrational spectra and force constant calculations of adsorption states of 2-propanol on TiO_2 surfaces
J. Weiss and V. Hoffmann (Tübingen, F.R.G.) 441

Excited complex formation between anthracene and *N,N*-diethylaniline on the silica surface
O.I. Yakimova, A.M. Eremenko and A.A. Chuiko (Kiev, U.S.S.R.) 447
Photoprotolytic interaction of acridine molecules with SiO_2 active centers
N.P. Smirnova, S.A. Samchuk, A.M. Yeremenko and A.A. Chuiko (Kiev, U.S.S.R.) 453

Author index 459
Subject index 462